

Characterization of Patients with Lung Cancer Not Meeting Lung Cancer Screening Criteria: Identifying a High-Risk Group Beyond Tobacco Smoking

JC. Laguna^{1,2}, A. Moll¹, M. Jiménez², J. Mena-Guerrero², M. Potrony³, L. Moreno⁴, E. Grau⁴, T. Gorria^{1,2}, M. Ferriol-Galmés^{2,5}, M. Liu², O. Castillo², N. Reguart^{1,2}, A. Arcocha¹, X. Solé^{2,3}, JA Puig-Butillé^{2,3}, C. Teixidó^{2,3,6}, T. Ramón y Cajal^{1,2}, L. Mezquita^{1,2}

1. Medical Oncology Department, Hospital Clinic Barcelona, Spain; 2. Translational Genomics and Targeted Therapeutics in Solid Tumors, IDIBAPS, Spain; 3. Department of Biochemistry & Molecular Genetics, Molecular Biology CORE laboratory, Hospital Clinic de Barcelona, Barcelona, Spain; 4. Gastroenterology Department, Hospital Clinic Barcelona, Spain; 5. Department of Computer Architecture, Universidad Politécnica de Cataluña, Barcelona, Spain; 6. Pathology Department, Hospital Clinic de Barcelona, Spain

laguna@clinic.cat

@jc_laguna_



Poster N#P3.01.46

BACKGROUND


- Lung cancer screening (LCS) programmes, based mainly on age and smoking history, have demonstrated a significant reduction in lung cancer (LC) mortality, prompting their gradual adoption¹⁻²
- However, **up to 40% of LC cases occurred in patients LCS-ineligible** according current LCS criteria.
- Susceptibility factors** reflected by family and personal history of cancer (FHC, PHC), could help to identify a **high-risk population**, not considered currently in LCS programmes.

OBJECTIVE:

To characterize the clinical and tumor profiles, as well as familial and individual susceptibility in patients with NSCLC who didn't meet LCS criteria

METHODS

- Prospective study** of 350 patients with **all-stages NSCLC** treated at Hospital Clinic of Barcelona (Spain) between Oct/2020 to Jan/2025.
- Patients had a **personal interview** to collect personal and family history of cancer (three-generation pedigree), demographic data, and exposure to environmental and occupational carcinogens. Clinical and molecular data was collected from medical records.
- Patients were classified as **LCS-eligible or ineligible at the time of NSCLC diagnosis** according to Nelson criteria [who had quit ≤ 10 years ago] who had smoked (50-74 years old (yo) current or former smokers >15 cigarettes a day for >25 years or >10 cigarettes a day for >30 years).
- The patient/tumor/family profile was analyzed in each LCS-ineligible group and compared with LCS-eligible patients.



Study population

Baseline characteristics N=350 (%)		
Age	Median, range	65 (30-88)
	Male	201 (57)
Gender	Female	149 (43)
Smoking status	Ever-smokers	289 (83)
	Non-smokers	61 (17)
Histology	Adenocarcinoma	255 (72)
	Squamous	51 (15)
	Others	44 (13)
	Stage	
Stage	Stage IV	187 (54)
	Stage I-III	161 (46)
PDL1 status	$\leq 1\%$	117 (33)
	1-49%	96 (27)
	$\geq 50\%$	91 (26)
	No data	46 (13)
Molecular status	Driver*	242 (74)
	No Driver	98 (28)
	Missing	12 (3)

Table 1. Baseline characteristics




Figure 1. Distribution of the study population by molecular status.

*Driver population was defined as somatic mutations (m) in EGFR/KRAS/BRAF/MET/ERBB2, fusions (f) in ALK/ROS1/RET/NTRK1-3, amplifications (a) in ERBB2/MET

RESULTS

Family & Personal History of Cancer in the LCS-inelegible population

Family History of Cancer

The prevalence of 1st-degree affected relatives was similar in both groups.




Figure 3. Proportion of patients with FHC in 1st-degree relatives by subgroup.

* By subgroup, the LCS-ineligible group due to non-smoking status showed the highest rate of FHC (82%), particularly colorectal (20%), breast (18%), and lung (12%) cancers.

Personal History of Cancer

- PHC was most common among LCS-inelegible patients





Figure 4. Proportion of patients with PHC in LCS-eligible and ineligible patients.

Figure 5. Cancer cases in patients from the LCS-inelegible group by sex. Haemato: Hamelatalogic; H&N: Head and neck; U: urinary

LCS-inelegible population



Out of 350 patients, **217 (62%) didn't meet LCS criteria** due to:

- Young age at NSCLC diagnosis (17%)
- Old age at NSCLC diagnosis (6%)
- Never-smoking status (22%)
- Low tobacco exposure (54%)




Figure 2. Flowchart of the study. LCS: Lung cancer screening; pop: population

Clinical & Molecular profile in LCS-inelegible population

A different profile was observed between LCS-ineligible and LCS-eligible populations, with a higher prevalence of female, lung adenocarcinoma, and driver alterations in the LCS-inelegible patients ($p<0.001$).




Figure 6. Radar plot with clinical/molecular profile in LCS-eligible and LCS-ineligible populations. LUAD: Lung adenocarcinoma; FHC: family history of cancer; PHC: personal history of cancer; yo: years old

CONCLUSIONS

- LCS current criteria exclude a significant proportion of individuals that will develop NSCLC.
- FHC may help to identify a high risk-population in LCS-ineligible group, particularly in non-smokers which may benefit of LCS.
- A deeper characterization of this population is currently being developed in a wider cohort (LAPIN study; EU4H-2024-PJ-04)

REFERENCES:

- Aberle et al. NEJM. 2011
- De Koning et al. NEJM 2020.
- Nandakumar et al. Eur Respir J. 2023

DISCLOSURES:

JCL has no disclosures to declare

FUNDING: JCL received support from Contractes Clínic de Recerca "Emili Letang-Josep Font" 2023; Hospital Clínic Barcelona, 2023; LM received support from EU4H-2024-PJ-04 and Beca SEOM-ASTRAZENECA para Proyectos de Investigación para Grupo Emergente 2022