

3056: *KRAS* Mutations Shape an Oxidative, Immune-Cold Tumor microenvironment Driving Chemo-immunotherapy Resistance in advanced NSCLC

IASLC

V. Seguí¹, R. Herrero¹, D. González-Cambor^{1,2}, J.A. Carbonell-Asins³, M. Cabeza-Segura¹, J. Martín-Arana¹, F. Gimeno-Valiente¹, I. Blasco¹, B. Iglesias¹, M. García-Bartolomé¹, N. Grimalt¹, J. Guijarro⁴, V. Schröer⁴, A. Compañ⁵, J. Franco-Serrano⁶, N. Tarazona^{1,2}, A. Cervantes^{1,2}, A. Insa¹, P. Martín-Martorell¹, J. Castillo⁷, V. Gambardella^{1,2}

¹ Oncology Department. Hospital Clínico Universitario. INCLIVA Biomedical Research Institute, Valencia/ES. ² CIBERONC. Instituto de Salud Carlos III, Madrid/ES. ³ Biostatistics Unit. INCLIVA Biomedical Research Institute, Valencia/ES. ⁴ Interventional Radiology Department. Hospital Clínico Universitario, Valencia/ES. ⁵ Pathology Department. Hospital Clínico Universitario, Valencia/ES. ⁶ Pneumology Department. Hospital Clínico Universitario, Valencia/ES. ⁷ University of Valencia, Valencia/ES

Correspondence: vic.seg.manz@gmail.com // valen.gambardella@gmail.com

INTRODUCTION

CONCLUSION

- Despite improved outcomes with checkpoint inhibitors (CPI) and Precision Medicine approaches, **NSCLC** is still the **leading cause of death by cancer**.
- ~ **30% of NSCLC** harbor **KRAS** mutations (G12C the most common and the only one with approved molecular-matched therapy).
- Beyond genomic alterations, the **tumor microenvironment** has emerged as a **critical determinant of treatment resistance**.
- Oxidative stress** has been shown to **modulate CPI efficacy** through altered PD-L1 expression and immune cell infiltration but the the precise **immunometabolic interplay** within the TME of *KRAS*-mutated tumors **remains to be elucidated**.

Our data suggest that **KRAS mutations** in NSCLC **shape a metabolically reprogrammed and immunosuppressive TME** that may drive resistance to ChT-IO.

Targeting **oxidative stress and immunomodulatory pathways**—such as those involving **EDN1**—may represent **promising therapeutic strategies** to overcome resistance and improve outcomes in *KRAS*-driven NSCLC.

RESULTS

Patients with **KRAS-mutant tumors** had significantly **shorter PFS** (p 0.006) (Fig 1). Lower *CD274* expression was associated with worse prognosis (p 0.005) and was balanced between groups (p 0.355). Conversely, a **KEAPness signature** assessment (Scalera et al CCR 2024) **replicates survival curves**, highlighting the importance of this pathway (Fig 2). Comparing the *KRAS*-mutant vs the *KRAS*-wt samples, there were no differences in the KEAPness signature expression (Fig 3A), but **patients with poor response** to treatment showed **increased expression of NFE2L2** gene (Fig 3B). Transcriptomic analysis underlined 83 differentially expressed genes between *KRAS*-mutant and *KRAS*-wt tumors, with 11 showing |log2FC| > 1 (Fig 4). **GSEA** demonstrated a **marked downregulation of immune-related pathways**—including antigen processing, interferon signaling, and T cell-mediated cytotoxicity—in *KRAS*-mutant tumors (Fig 5). The **CIBERSORT immune deconvolution** highlights a **general downregulation of the immune components** in the *KRAS*-mutant samples (Fig 6). Overexpressed in *KRAS*-mutant tumors (log2FC = 1.16; p 0.026), **EDN1** was the gene **most negatively correlated with CD8+ T cell infiltration** (Fig 7), with **known roles in ROS metabolism and TGFβ-mediated immunosuppression**. EDN1 is **differentially expressed** (p 0.026) (Fig 7A) and its **higher expression** is related with **lower survival** (p 0,001) (Fig 7B).

METHODS

Prospective collection of samples from **47 patients**:

- Advanced NSCLC
- Treatment naïve, candidates to receive ChT-IO
- Included **KRAS-mut** and **KRAS-wt** cases (oncogene-addicted cases excluded)

RNA extracted from samples with ≥30% tumor cellularity and processed using the TruSeq RNA Access Kit (Illumina), followed by paired-end sequencing on the NovaSeq 6000 platform

Transcriptomic data pre processed via nf-core/rnaseq. Differential gene expression analysis conducted using DESeq2, with downstream gene set enrichment analysis (GSEA). Immune landscape inference using CIBERSORT and ESTIMATE algorithms.

Associations between gene expression, immune composition, and clinical outcomes were evaluated using Spearman correlation, Cox multivariate modeling, and Kaplan-Meier analysis.